

COMPANY ANNOUNCEMENT

Vi-Jon, LLC Expands Voluntary Nationwide Recall of All Flavors and Lots Within Expiry of Magnesium Citrate Saline Laxative Oral Solution Due to Microbial Contamination

When a company announces a recall, market withdrawal, or safety alert, the FDA posts the company's announcement as a public service. FDA does not endorse either the product or the company.

Read Announcement

View Product Photos

More Recalls, Market Withdrawals, & Safety Alerts (/safety/recalls-market-withdrawals-safety-alerts)

Summary

Company Announcement Date:

July 25, 2022

FDA Publish Date:

July 26, 2022

Product Type:

Drugs

Reason for Announcement:

Potential Gluconacetobacter liquefaciens contamination

Company Name:

Vi-Jon, LLC

Brand Name:

Multiple Brands

Product Description:

Magnesium Citrate Saline Laxative Oral Solution

Company Announcement

FOR IMMEDIATE RELEASE – July 25, 2022 – Smyrna, TN, Vi-Jon, LLC is expanding its voluntary recall to include all lots of all flavors of Magnesium Citrate Saline Laxative Oral Solution within expiry to the consumer level. This expansion includes all lots of Cherry Flavor and Grape Flavor of Magnesium Citrate Saline Laxative Oral Solution, 10 FL OZ (296 mL) within expiry. On July 14, 2022, Vi-Jon, LLC recalled all lots of Magnesium Citrate Saline Laxative Oral Solution Lemon Flavor, 10 FL OZ (296 mL) within expiry. The recall was initiated after Vi-Jon, LLCs third party microbial testing identified the presence of *Gluconacetobacter liquefaciens*.

Risk Statement: Immunocompromised patients, who consume this product, may be at increased risk for invasive infections caused by *Gluconacetobacter liquefaciens* that could lead to serious, life-threatening adverse health consequences. To date, Vi-Jon, LLC is aware of **3 (three)** reports of serious adverse reactions potentially related to this recall. Vi-Jon, LLC is in the process of investigating these reports. The product is used for relief of occasional constipation (irregularity) and generally produces bowel movement in ½ to 6 hours. The product is packaged in a 10 oz clear round plastic bottle.

In addition to the lemon flavor, the recall now includes the Cherry flavor and Grape flavor noted below in the table in bold print.

Affected Brand	NDC #	UPC #
BEST CHOICE 100Z LEMON MAG CITRATE	63941-533-38	070038200499
BEST CHOICE 100Z CHERRY CITRATE	63941-516-38	070038587903
BEST CHOICE 100Z GRAPE CITRATE	63941-162-38	070038662204
CARE ONE 100Z LEMON MAG CIT	72476-001-38	341520313226
CARE ONE 100Z CHERRY CIT	72476-002-38	341520000553
CARIBA 100Z LEMON MAG CITRATE	67860-166-38	646702057012
CRUZ BLANC 100Z LEMON MAG CIT	N/A	308697403082
CVS 100Z LEMON MAG CIT	63868-929-38	050428335178
CVS 100Z LEMON MAG CIT	69842-983-38	050428305942
CVS 100Z CHERRY CITRATE	69842-647-38	050428297339
CVS 100Z CHERRY CITRATE	69842-647-38	00050428285152
CVS 100Z CLR GRAPE CITRATE	69842-763-38	050428307458
CVS 100Z CLR GRAPE CITRATE	69842-763-38	00050428325032
DISCOUNT DRUG MART 100Z LEMON MAG CITRATE	53943-166-38	093351028205
EQUALINE 100Z LEMON MAG CIT	41163-709-38	041163500679
EQUALINE 100Z CHERRY CITRATE	41163-769-38	041163500686
EQUATE 100Z LEMON MAG CIT SRP	49035-506-38	681131287142
EQUATE 100Z CHERRY CIT SRP	49035-593-38	681131287166
EQUATE 100Z GRAPE MAG CIT SRP	49035-592-38	681131287159
EXCHANGE SELECT 100Z LEMON MAG CIT	55301-166-38	614299404205
FAMILY WELLNESS 100Z LEMON CITRATE	55319-666-38	032251580826
FAM WELLNS 100Z CHERRY CITRATE	55319-164-38	032251577888
GOOD SENSE 100Z LEMON MAG CIT	50804-166-38	846036007374
GOOD SENSE 100Z CHERRY CITRATE	50804-164-38	846036007398

Affected Brand	NDC #	UPC #
HARRIS TEETER 100Z LEMON MAG CITRATE	72036-002-38	072036726124
HEB 100Z LEMON MAG CITRATE	37808-769-38	041220510863
HEB 100Z CHERRY CITRATE	37808-673-38	00041220510870
HEB 100Z GRAPE MAG CITRATE	37808-695-38	00041220510887
HEALTH MART 100Z LEMON MAG CIT	62011-0380-1	052569142158
HEALTH MART 100Z CHERRY CIT	62011-0381-1	052569142165
KROGER 100Z LEMON MAG CITRATE	30142-899-38	041260001826
KROGER 100Z GRAPE CITRATE	30142-806-38	041260008719
LEADER 100Z LEMON MAG CIT	70000-0424-1	096295135541
LEADER 100Z CHERRY CIT	70000-0575-1	096295141061
LEADER 100Z GRAPE MAG CIT	70000-0576-1	096295141054
MAJOR 100Z LEMON MAG CITRATE	0904-6787-44	309046787440
MEIJER 100Z LEMON MAG CIT	41250-708-38	713733459457
MEIJER 100Z CHERRY CITRATE	41250-769-38	713733459440
PREMIER VALUE 100Z LOW SOD LEM CIT	68016-696-38	840986035302
PREMIER VALUE 100Z CHERRY CIT	68016-701-38	840986035296
PUBLIX 100Z LEMON MAG CIT	56062-266-38	041415506732
PUBLIX 100Z CHERRY CITRATE	56062-264-38	041415505735
QUALITY CHOICE 100Z LEMON MAG CIT	63868-929-38	635515901254
QUALITY CHOICE 100Z CHERRY CITRATE	63868-018-38	635515901117
REXALL 100Z LEMON MAG CITRATE	55910-183-38	072785134188
REXALL 100Z CHERRY CIT	55910-961-38	072785134164
REXALL 100Z GRAPE MAG CIT	55910-615-38	072785134171
RITE AID 100Z LEMON CITRATE	11822-4330-2	011822433006
RITE AID 100Z CHERRY CITRATE	11822-4303-2	011822433037
SIGNATURE CARE 100Z LEMON MAG CIT	21130-709-38	321130779155
SIGNATURE CARE 100Z CHERRY CIT	21130-165-38	321130789710
SOUND BODY 100Z LEMON MAG CIT	50594-166-38	072785114791
SUNMARK 100Z LEMON MAG CIT	70677-0051-1	010939908445
SUNMARK 100Z CHERRY CIT	70677-0053-1	010939910448
SWAN 100Z LEMON MAG CITRATE	0869-0166-38	072785134058
SWAN 100Z CHRY CITRATE	0869-0164-38	308690693381
TOPCARE 100Z LEMON MAG CITRATE	36800-709-38	036800455290
TOPCARE 100Z CHERRY CIT	36800-164-38	036800455306
UP&UP 100Z LEMON MAG CIT	11673-708-38	072785128835
UP&UP 100Z LEMON MAG CIT	11673-666-38	072785128835
WALGREENS 100Z LEMON MAG CIT	0363-8166-38	311917201603
WALGREENS 100Z CHERRY CIT	0363-8164-38	311917201580
WALGREENS 100Z GRAPE MAG CIT	0363-7162-38	311917201597

The product was distributed Nationwide to wholesale and retail outlets. Vi-Jon, LLC is continuing their investigation into the cause of the problem.

Vi-Jon, LLC is notifying its customers by phone and email and is arranging for return or destruction of all recalled product. Consumers that have this recalled product should stop using and return any remaining product to the place of purchase.

Consumers with questions regarding this recall can contact Vi-Jon, LLC by e-mail (Recalls@Vijon.com (<mailto:Recalls@Vijon.com>)) Monday-Friday, from 7:30 am to 4:30 pm, Central Time. Consumers should contact their physician or healthcare provider if they have experienced any problems that may be related to taking or using this drug product www.vijon.com (<http://www.vijon.com/>) [🔗](http://www.fda.gov/about-fda/website-policies/website-disclaimer) (<http://www.fda.gov/about-fda/website-policies/website-disclaimer>).

Adverse reactions or quality problems experienced with the use of this product may be reported to the FDA's MedWatch Adverse Event Reporting program either online, by regular mail or by fax.

- Complete and submit the report [Online](#) (</safety/medwatch-fda-safety-information-and-adverse-event-reporting-program/reporting-serious-problems-fda>)
- Regular Mail or Fax: [Download form](#) (</safety/medical-product-safety-information/medwatch-forms-fda-safety-reporting>) or call 1- 800-332-1088 to request a reporting form, then complete and return to the address on the pre-addressed form, or submit by fax to 1-800-FDA-0178

This recall is being conducted with the knowledge of the U.S. Food and Drug Administration.

[Link to 1st Expansion Press Release](#) (</safety/recalls-market-withdrawals-safety-alerts/vi-jon-llc-expands-voluntary-nationwide-recall-magnesium-citrate-saline-laxative-oral-solution-lemon>).

[Link to Initial Press Release](#) (</safety/recalls-market-withdrawals-safety-alerts/vi-jon-llc-issues-voluntary-nationwide-recall-cvs-magnesium-citrate-saline-laxative-oral-solution>).

Company Contact Information

Consumers:

Vi-Jon, LLC

✉ Recalls@Vijon.com (<mailto:Recalls@Vijon.com>)

Media:

Mr. Joseph Meehan, Chief Sales and Marketing Officer

☎ 615-208-2441

Product Photos



Drug Facts Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g Saline laxative	Drug Facts (continued) or contact a Poison Control Center right away. Store at temperatures between 40° and 80° F (4° and 30° C)
Uses ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in 1/2 to 6 hours	Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Warnings Ask a doctor before use if you have: ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help.	Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water PROUDLY DISTRIBUTED BY: VALU MERCHANDISERS, CO. 5000 KANSAS AVE KANSAS CITY, KS 66106 L00194825A




Drug Facts Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g Saline laxative	Drug Facts (continued) or contact a Poison Control Center right away. Store at temperatures between 40° and 80° F (4° and 30° C)
Uses ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in 1/2 to 6 hours	Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Warnings Ask a doctor before use if you have: ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help.	Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water PROUDLY DISTRIBUTED BY: VALU MERCHANDISERS, CO. 5000 KANSAS AVE KANSAS CITY, KS 66106 L00194825A



Drug Facts Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g Saline laxative	Drug Facts (continued) or contact a Poison Control Center right away. Store at temperatures between 40° and 80° F (4° and 30° C)
Uses ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in 1/2 to 6 hours	Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Warnings Ask a doctor before use if you have: ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help.	Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water Questions? Call 1-877-846-9949

DISTRIBUTED BY: FOODHOLD U.S.A., LLC, LANDOVER, MD 20785
 For product questions or concerns, contact us at 1-877-846-9949
 © 2018 Ahold Delhaize Licensing Sarl • Quality guaranteed or your money back.





Magnesium Citrate

SALINE LAXATIVE
Oral solution
Non-pasteurized

DYE
FREE



Cherry Flavor
Alcohol content 0.0250%
(from flavoring)

10 FL OZ (296 mL)

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKRING

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.

■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.
Store at temperatures between 40° and 80° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age ask a doctor

Other information

■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

L00175885A

#231183

Distributed by: CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02895
© 2016 CVS Pharmacy
CVS.com 1-800-SHOP-CVS
V-16463



Magnesium Citrate

SALINE LAXATIVE
Oral solution
Non-pasteurized

DYE
FREE



Grape Flavor
Alcohol content 0.0013%
(from flavoring)

10 FL OZ (296 mL)

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKRING

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.

■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.
Store at temperatures between 40° and 80° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age ask a doctor

Other information

■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

L00175755A

#789407

Distributed by: CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02895
© 2016 CVS Pharmacy
CVS.com 1-800-SHOP-CVS
V-16463



NDC 41163-769-38

EQUALINE®

magnesium citrate

saline laxative
oral solution

• saccharin free
• non-pasteurized

ALCOHOL CONTENT 0.0250% (FROM FLAVORING)

DYE FREE

cherry flavor



10 FL OZ (296 mL)

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKRING

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.

■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.
Store at temperatures between 40° and 80° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age ask a doctor

Other information

■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions? 1-877-932-7948



L00175905A

Distributed by: SUPERVALU INC.
Eden Prairie, MN 55344 USA
877-932-7948
supervaluprivatelabels.com

Magnesium Citrate
Saline Laxative

• Relief of occasional constipation
• For use in bowel cleansing systems for surgery or exam preparation
• Non-pasteurized

Alcohol content 0.0250% (from flavoring)

10 FL OZ (296mL)

Cherry Flavor

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g.....Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in 1/2 to 6 hours

Warnings
Ask a doctor before use if you have:
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are:
■ taking any other drug.
■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.
Store at temperatures between 40° and 86°F (8° and 30°C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information
■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions? 1-888-287-1915

Do not use if tamper evident twist-off cap is missing, broken or separated from neck ring.

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915

DISTRIBUTED BY: Wal-Mart Inc., Bentonville, AR 72716

6 81131 28716 6

L0018321SA 539627

Magnesium Citrate
Saline Laxative

• Relief of occasional constipation
• For use in bowel cleansing systems for surgery or exam preparation

Alcohol content 0.0013% (from flavoring)

10 FL OZ (296mL)

Grape Flavor

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g.....Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in 1/2 to 6 hours

Warnings
Ask a doctor before use if you have:
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are:
■ taking any other drug.
■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.
Store at temperatures between 40° and 86°F (8° and 30°C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information
■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions? 1-888-287-1915

Do not use if tamper evident twist-off cap is missing, broken or separated from neck ring.

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915

DISTRIBUTED BY: Wal-Mart Inc., Bentonville, AR 72716

6 81131 28715 9

L0018320SA 539627

Magnesium Citrate
Saline laxative

Alcohol content 0.0250% (from flavoring)

10 FL OZ (295 mL)

CHERRY

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g.....Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in 1/2 to 6 hours

Warnings
Ask a doctor before use if you have:
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are:
■ taking any other drug.
■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.
Store at temperatures between 40° and 86°F (8° and 30°C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information
■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

NOT 100% SATISFIED?
Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

Do not use if tamper evident twist-off cap is missing, broken or separated from neck ring.

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915

DISTRIBUTED BY: MIDWOOD BRANDS, LLC 500 VOLVO PARKWAY CHESAPEAKE, VA 23320

0 32251 57788 8

L00189488A SKU 902283

NDC 50804-164-38

GOODSENSE

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKING

Magnesium Citrate

Saline Laxative

Oral Solution

• Non-pasteurized • Saccharin free

Cherry Flavor

Alcohol content 0.0250% (from flavoring)

10 FL OZ (296 mL)

100% SATISFACTION GUARANTEED

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.
■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.

Store at temperatures between 40° and 86° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information

■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions? Call 1-888-593-0593

0 65200 95094
0 65200 10739
8
Distributed by:
Geiss, Destin & Dunn, Inc.
Peachtree City, GA 30269
www.valuelabels.com
L00175949A

H-E-B

Magnesium Citrate

Oral Solution

Saline Laxative

• Relief of:
Occasional Constipation

Sugar & Dye Free

Gluten Free

Alcohol content 0.0250% (from flavoring)

Cherry Flavor

10 FL OZ (296 mL)

Do Not Use If Tamper Evident Twist-off Cap Is Missing, Broken or Separated From Necking

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.
■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.

Store at temperatures between 40° and 86° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information

■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions? Call 1-888-593-0593

0 18015 02214
0 18015 02214
4 1220 51087
0
MADE WITH PRIDE & CARE FOR H-E-B®
SAN ANTONIO, TX 78204
L00175835A
9761-1808

H-E-B

Magnesium Citrate

Oral Solution

Saline Laxative

• Relief of:
Occasional Constipation

Sugar & Dye Free

Gluten Free

Alcohol content 0.0013% (from flavoring)

Grape Flavor

10 FL OZ (296 mL)

Do Not Use If Tamper Evident Twist-off Cap Is Missing, Broken or Separated From Necking

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.
■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.

Store at temperatures between 40° and 86° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information

■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions? Call 1-888-593-0593

0 88015 02214
0 88015 02214
4 1220 51088
7
MADE WITH PRIDE & CARE FOR H-E-B®
SAN ANTONIO, TX 78204
L00175745A
9761-1808

HealthMart.

NDC 62011-0361-1

Magnesium Citrate

Saline Laxative and Oral Solution

Saccharin free
Break resistant
Very low sodium
BPA free

CHERRY FLAVORED
Alcohol content 0.0250% (from flavoring)

10 FL OZ (296 mL)

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKRING

Non-pasteurized

Drug Facts

Active ingredient (in each fl oz) Magnesium citrate 1.745 g

Purpose Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in ½ to 6 hours

Warnings
Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.
■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of

Drug Facts (continued)
overdose, get medical help or contact a Poison Control Center right away.
Store at temperatures between 40° and 80° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information
■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

McKesson
Distributed by McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee
healthmart.com

0 52569 14216 5

VS61521001

Kroger

OUR PHARMACIST RECOMMENDS

Magnesium Citrate

SALINE LAXATIVE

Alcohol content 0.0013% (from flavoring)

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKRING

Grape Flavor

10 FL OZ (296 mL)

Drug Facts

Active ingredient (in each fl oz) Magnesium citrate 1.745 g

Purpose Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in ½ to 6 hours

Warnings
Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.
■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help

Drug Facts (continued)
or contact a Poison Control Center right away.
Store at temperatures between 40° and 80° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information
■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions? Call 1-800-632-6900

DISTRIBUTED BY THE KROGER CO.
CINCINNATI, OHIO 45202
800-632-6900 | www.kroger.com

0 41260 00871 9

L0018960SA

LEADER²

NDC 70000-0575-1

Sugar-Free Magnesium Citrate

Saline Laxative

Cherry Flavor

Non-Pasteurized
Break Resistant
BPA-Free

100% Money Back Guarantee

10 FL OZ (296 mL)

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKRING

Drug Facts

Active ingredient (in each fl oz) Magnesium citrate 1.745 g

Purpose Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in ½ to 6 hours

Warnings
Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.
■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help

Drug Facts (continued)
or contact a Poison Control Center right away.
Store at temperatures between 40° and 80° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information
■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions? Call 1-888-593-0593

DISTRIBUTED BY CARDINAL HEALTH
COLUMBUS, OHIO 43007
www.mylaxative.com
1-800-200-6393
©2017 Cardinal Health
CIN 5725452
REV. 10/21

0 9629514106 1

L0019884SA

LEADER²

NDC 70000-0576-1

**Sugar-Free
Magnesium
Citrate**
Saline Laxative

**Grape
Flavor**

Non-Pasteurized
Break Resistant
BPA-Free

**100% Money
Back Guarantee**

10 FL OZ (296 mL)

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKING

Drug Facts	Drug Facts (continued)
Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g..... Saline laxative	or contact a Poison Control Center right away. Store at temperatures between 40° and 86° F (4° and 30° C)
Uses ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in ½ to 6 hours	Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Warnings Ask a doctor before use if you have ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are ■ taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work. Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
	Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg
	Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water
	Questions? Call 1-888-593-0593

DISTRIBUTED BY
CARDINAL HEALTH
DUBLIN, OHIO 43017
www.myleader.com
1-800-200-5315
©2020 Cardinal Health
CIN 5725445
REV. 10/21



0 92625 14105

L0019883SA

NDC 41250-769-38

Do not use if tamper evident twist-off cap is missing, broken or separated from necking

meijer

**magnesium
citrate**
Sugar Free | Low Sodium
Saline Laxative

Relief of Occasional Constipation

Contains no ingredient made from gluten-containing grain (wheat, barley or rye)

CHERRY
Alcohol content 0.0250% (from flavoring)

10 FL OZ (296 mL)

Drug Facts	Drug Facts (continued)
Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g..... Saline laxative	or contact a Poison Control Center right away. Store at temperatures between 40° and 86° F (4° and 30° C)
Uses ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in ½ to 6 hours	Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Warnings Ask a doctor before use if you have ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are ■ taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work. Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
	Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg
	Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water
	Questions? 1-888-593-0593

DIST. BY MEIJER
DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com



7 13731 45044

L00175875A

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKING

**Premier
Value**

**Magnesium
Citrate**
saline laxative
oral solution

Non-pasteurized
Saccharin free
BPA free
Dye Free

Cherry Flavor
Alcohol content 0.0250% (from flavoring)

10 FL OZ (296 mL)

Drug Facts	Drug Facts (continued)
Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g..... Saline laxative	Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Use ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in ½ to 6 hours	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
Warnings Ask a doctor before use if you have ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are ■ taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work. Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Store at temperatures between 40° and 86° F (4° and 30° C)	Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg
	Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water
	Questions? Call 1-888-593-0593

L00175965A



8 40986 03529 6

Distributed by:
Pharmacy Value Alliance LLC
407 East Lancaster Avenue,
Wayne, PA 19087

<h3>Drug Facts</h3> <p>Active ingredient (in each 1/2 oz) Purpose Magnesium chloride 1.745 g Laxative</p>	<h3>Drug Facts (continued)</h3> <p>or contact a Poison Control Center right away. Store at temperatures between 49° and 89° F (5° and 30° C).</p>
<p>Uses • for relief of occasional constipation (irregularity). • Generally produces bowel movement in 1/4 to 6 hours.</p>	<h3>Directions • shake well before using • drink a full glass (8 ounces) of liquid with each dose • may be taken as a single daily dose or in divided doses</h3>
<h3>Warnings</h3> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • kidney disease • a magnesium restricted diet • abdominal pain, nausea, or vomiting • noticed a sudden change in bowel habits that persists over a period of 2 weeks • already used a laxative for a period longer than 1 week <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> • taking any other drug. • Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work. <p>Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.</p>	<p>adults and children 12 years of age and over 6.5 to 10 1/2 oz maximum 10 1/2 oz in 24 hours</p> <p>children 6 to under 12 years of age 3 to 6 oz maximum 7 1/2 oz in 24 hours</p> <p>children 2 to under 6 years of age 2 to 3 1/2 oz maximum 3 1/2 oz in 24 hours</p> <p>children under 2 years of age ask a doctor</p>
	<h3>Other information</h3> <ul style="list-style-type: none"> • each 1/2 oz contains: magnesium 290 mg • each 1/2 oz contains sodium 1 mg <h3>Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water</h3>

52505 SL414 41415 50573

<h2>Drug Facts</h2> <p>Active ingredient (in each fl oz) Purpose Magnesium citrate 1.74 g Saline laxative</p> <p>Uses ■ for relief of occasional constipation (regularity). ■ Generally produces bowel movement in 7½ to 6 hours</p> <h2>Warnings</h2> <p>Ask a doctor before use if you have ■ kidney disease or a magnesium restricted diet. ■ abdominal pain, nausea, or vomiting. ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks. ■ already used a laxative for a period longer than 1 week</p> <p>Ask a doctor or pharmacist before use if you are ■ taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.</p>	<h2>Drug Facts (continued)</h2> <p>or contact a Poison Control Center right away. Store at temperatures between 40° and 80° F (4° and 30° C).</p> <h2>Directions</h2> <p>■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses</p> <table border="1"> <tr> <td>adults and children 12 years of age and over</td> <td>6.5 to 10 fl oz maximum, 10 fl oz in 24 hours</td> </tr> <tr> <td>children 6 to under 12 years of age</td> <td>3 to 7 fl oz or maximum 7 fl oz in 24 hours</td> </tr> <tr> <td>children 2 to under 6 years of age</td> <td>2 to 3 fl oz or maximum 3 fl oz in 24 hours</td> </tr> <tr> <td>children under 2 years of age</td> <td>ask a doctor</td> </tr> </table>	adults and children 12 years of age and over	6.5 to 10 fl oz maximum, 10 fl oz in 24 hours	children 6 to under 12 years of age	3 to 7 fl oz or maximum 7 fl oz in 24 hours	children 2 to under 6 years of age	2 to 3 fl oz or maximum 3 fl oz in 24 hours	children under 2 years of age	ask a doctor
adults and children 12 years of age and over	6.5 to 10 fl oz maximum, 10 fl oz in 24 hours								
children 6 to under 12 years of age	3 to 7 fl oz or maximum 7 fl oz in 24 hours								
children 2 to under 6 years of age	2 to 3 fl oz or maximum 3 fl oz in 24 hours								
children under 2 years of age	ask a doctor								
<h2>Other information</h2> <p>■ each fl oz contains: magnesium 280 mg ■ each fl oz contains: sodium 1 mg</p> <h2>Inactive ingredients</h2> <p>benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water, yellow 6</p>									

6
35515 90111
7

Drug Facts

Active Ingredient (in each fl oz) Purpose
Magnesium chloride 1.745 g _____ Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in 7½ to 6 hours

Warnings
Ask a doctor before use if you have
■ kidney disease or a magnesium-patched diet
■ additional pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other medicines
■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help.

Drug Facts (continued)

or contact a Poison Control Center right away.
Store at temperatures between 49° and 86° F (5° and 30° C).
Keep out of reach of children.

Directions ■ shake well before using ■ drink full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10.0 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information
■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzocaine, citric acid, disodium EDTA, flavor, sucralose, water.

Questions? Call 1-866-4-REXALL.

Visit us at: Revall.com or call 1-866-4-REXALL 7

DISTRIBUTED BY DOLGENCORP, LLC

100 MISSION RIDGE

A0637

91751 58222 0

Since 1903 Rexall® is a registered trademark of Rexall Sundown, Inc.

Since 1903
Rexall[®]

Magnesium Citrate Saline Laxative

- Oral solution
- Relief of occasional constipation

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECK RING
Alcohol Content 0.0013% (from flavoring)

10 FL OZ (296 mL)

Grape Flavored

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g.....Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.
Store at temperatures between 40° and 80° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information

■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions? Call 1-866-4-REXALL

Since 1903 Rexall[®] is a registered trademark of Rexall Sundown, Inc.

Visit us at: Rexall.com or call 1-866-4-REXALL
DISTRIBUTED BY OLD EAST MAIN CO.
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072 L0017525B

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECK RING
NDC 11822-4303-2

MAGNESIUM CITRATE

SALINE LAXATIVE

Oral solution
Sugar-free

Alcohol content
0.0250%
(from flavoring)

CHERRY FLAVOR

10 FL OZ (296 mL)

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g.....Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.
Store at temperatures between 40° and 80° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information

■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.
DISTRIBUTED BY:
RATE HQ
30 HUNTER LANE
CHAP HILL, PA 17011
www.rtaxall.com
11822-4303-2

L0019362SA

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECK RING
NDC 11822-4303-2

MAGNESIUM CITRATE

SALINE LAXATIVE

Oral solution
Sugar-free

Alcohol content
0.0250%
(from flavoring)

CHERRY FLAVOR

10 FL OZ (296 mL)

Drug Facts

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 g.....Saline laxative

Uses ■ for relief of occasional constipation (irregularity).
■ Generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have
■ kidney disease ■ a magnesium restricted diet
■ abdominal pain, nausea, or vomiting
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ taking any other drug.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help

Drug Facts (continued)

or contact a Poison Control Center right away.
Store at temperatures between 40° and 80° F (4° and 30° C)

Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other information

■ each fl oz contains: magnesium 290 mg
■ each fl oz contains: sodium 1 mg

Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.
DISTRIBUTED BY:
RATE HQ
30 HUNTER LANE
CHAP HILL, PA 17011
www.rtaxall.com
11822-4303-2

L0019362SB

NDC 21130-165-38

Signature Care
Quality Guaranteed

Oral Solution
Magnesium Citrate
SALINE LAXATIVE
SACCHARIN FREE
Alcohol content 0.0250% (from flavoring)
CHERRY FLAVOR

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKING

10 FL OZ (296 mL)

Drug Facts Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g.....Saline laxative		Drug Facts (continued) Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses	
Use for relief of occasional constipation (irregularity). Generally produces bowel movement in ½ to 6 hours		adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours	
Warnings Ask a doctor before use if you have ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work. Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Store at temperatures between 46° and 86° F (8° and 30° C)		children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor	
Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg		Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water	

RD 18176 S2174

3 21130 78971 0

DISTRIBUTED BY BETTER LIVING BRANDS LLC
P.O. BOX 99, FLEXINGTON, CA 95051-0099
1-888-723-3929 www.betterlivingbrandsllc.com

L0017565SA

sunmark™ NDC 70677-0053-1

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECKING

magnesium citrate
Saline Laxative
Oral Solution
Alcohol content 0.0250% (from flavoring)
Non-pasteurized
Saccharin free
Break resistant
Very low sodium
BPA free
CHERRY FLAVOR

Distributed By McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee

10 FL OZ (296 mL)

Drug Facts Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g.....Saline laxative		Drug Facts (continued) or contact a Poison Control Center right away. Store at temperatures between 46° and 86° F (8° and 30° C)	
Use for relief of occasional constipation (irregularity). Generally produces bowel movement in ½ to 6 hours		Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses	
Warnings Ask a doctor before use if you have ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work. Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help		adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor	
Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg		Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water	

0 10939 91044 8

L0017577SA

NDC 0869-0164-38

2 swan.

VERY LOW SODIUM
Citroma
MAGNESIUM CITRATE

cherry
• Saline Laxative
• Oral Solution
Alcohol content 0.0250% (from flavoring)

Do not use if tamper evident twist-off cap is missing, broken or separated from necking

10 FL OZ (296 mL)

Drug Facts Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g.....Saline laxative		Drug Facts (continued) or contact a Poison Control Center right away. Store at temperatures between 46° and 86° F (8° and 30° C)	
Uses ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in ½ to 6 hours		Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses	
Warnings Ask a doctor before use if you have ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work. Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help		adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor	
Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg		Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water	

85369 06980 3

Distributed by: V-L-Jon, Inc.
One Swan Drive,
Smyrna, TN 37167

L0017726SA

+TopCare
health

NDC 36800-164-38

Magnesium Citrate

SALINE LAXATIVE ORAL SOLUTION

Alcohol content 0.0250% (from flavoring)

For Relief of Occasional Constipation

CHERRY FLAVOR


10 FL OZ (296 mL)

TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

OUR PHARMACISTS RECOMMEND

Drug Facts	Drug Facts (continued)
Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g.....Saline laxative	or contact a Poison Control Center right away. Store at temperatures between 49° and 68° F (8° and 33° C).
Uses ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in 1/2 to 6 hours	Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Warnings Ask a doctor before use if you have ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are ■ taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help	Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water
	Questions? Call 1-888-423-0139

DISTRIBUTED BY TOPCO ASSOCIATES LLC ELK GROVE VILLAGE, IL 60007
©TOPCO VJAO818 QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com
Visit here for more information: http://topbrnds.com/48zryu



Walgreens

NDC 0363-8164-38

Dye-Free Magnesium Citrate

SALINE LAXATIVE ORAL SOLUTION

SUGAR FREE

LOW SODIUM

Alcohol content 0.0250% (from flavoring)

10 FL OZ (296 mL)

TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

CHERRY FLAVOR

Drug Facts	Drug Facts (continued)
Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g.....Saline laxative	or contact a Poison Control Center right away. Store at temperatures between 49° and 68° F (8° and 33° C).
Uses ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in 1/2 to 6 hours	Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Warnings Ask a doctor before use if you have ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are ■ taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help	Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water
	Questions? Call 1-888-593-0593

ITEM 873223 W10100-0618-F L0017581SA ORG0518-F2 REV-0618



Walgreens

NDC 0363-8164-38

Dye-Free

Magnesium Citrate

SALINE LAXATIVE ORAL SOLUTION

Sugar Free

Low Sodium

Alcohol content 0.0250% (from flavoring)


10 FL OZ (296 mL)

TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

CHERRY FLAVOR

Drug Facts	Drug Facts (continued)
Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g.....Saline laxative	or contact a Poison Control Center right away. Store at temperatures between 49° and 68° F (8° and 33° C).
Uses ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in 1/2 to 6 hours	Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Warnings Ask a doctor before use if you have ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are ■ taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help	Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water
	Questions? Call 1-888-593-0593

Walgreens Pharmacist Recommended.
Our pharmacists recommend the Walgreens brand.
ITEM 873223 W10100-0821-F L0017581SB W3ORG0821-F



Walgreens NDC 0363-7162-38

Dye-Free

Magnesium Citrate

SALINE LAXATIVE ORAL SOLUTION

SUGAR FREE

LOW SODIUM

GRAPE FLAVOR

Alcohol content 0.0013% (from flavoring)

10 FL OZ (296 mL)

TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING

Drug Facts	Drug Facts (continued)
Active Ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g.....Saline laxative	or contact a Poison Control Center right away. Store at temperatures between 46° and 68° F (8° and 30° C).
Uses ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in 1½ to 6 hours	Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Warnings Ask a doctor before use if you have ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are ■ taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work. Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
	Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg
	Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water
	Questions? Call 1-888-593-0593

ITEM 873316 W10100-0618-F

L0017573SA

ORG0518-F2 REV-0618

3
191720159
7
DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2018 Walgreen Co.

Walgreens NDC 0363-7162-38

Dye-Free

Magnesium Citrate

SALINE LAXATIVE ORAL SOLUTION

Sugar Free

Low Sodium

Grape flavor

10 FL OZ (296 mL)

TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING

Drug Facts	Drug Facts (continued)
Active Ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g.....Saline laxative	or contact a Poison Control Center right away. Store at temperatures between 46° and 68° F (8° and 30° C).
Uses ■ for relief of occasional constipation (irregularity). ■ Generally produces bowel movement in 1½ to 6 hours	Directions ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Warnings Ask a doctor before use if you have ■ kidney disease ■ a magnesium restricted diet ■ abdominal pain, nausea, or vomiting ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ already used a laxative for a period longer than 1 week Ask a doctor or pharmacist before use if you are ■ taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work. Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
	Other information ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg
	Inactive ingredients benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water
	Questions? Call 1-888-593-0593

Walgreens Pharmacist Recommended.

Our pharmacists recommend the Walgreens brand.

ITEM 873316 W10100-0821-F

L0017573SB W30RG0821-F

3
191720159
7
DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.